

Institut für Lebensmittelsicherheit und -hygiene
der Vetsuisse-Fakultät Universität Zürich

Direktor: Prof. Dr. med. vet. Dr. h.c. Roger Stephan

**Shiga toxin-producing *Escherichia coli* (STEC) isolated from fecal samples of African
dromedary camels**

Inaugural-Dissertation

zur Erlangung der Doktorwürde der
Vetsuisse-Fakultät Universität Zürich

vorgelegt von

Melinda Baschera

Tierärztin
von Zürich

genehmigt auf Antrag von

Prof. Dr. med. vet. Dr. h.c. Roger Stephan, Referent

2020

Inhaltsverzeichnis	
Summary	4
Zusammenfassung	5
Shiga toxin-producing <i>Escherichia coli</i> (STEC) isolated from fecal samples of African dromedary camels	6
Abstract	7
Introduction	8
Material and Methods	10
Results and discussion	13
Conclusions	16
Nucleotide sequences	16
Disclosures	16
Acknowledgements	16
References	18
Danksagung	
Curriculum Vitae	

Summary

Shiga toxin producing *Escherichia coli* (STEC) cause gastrointestinal illnesses. To investigate the occurrence of STEC among Kenyan dromedaries, *E. coli* isolated from 163 dromedaries were screened for *stx1* and *stx2*. STEC strains were isolated and serotyped. Isolates were subjected to PCR for the subtyping of *stx* genes and for the detection of *eae* and *ehx*. Whole genome sequencing (WGS) was carried out to detect further virulence genes and to determine the multilocus sequence types (MLST). Antimicrobial resistance profiles were determined by disk diffusion.

STEC was isolated from 20 (12.3%) of the samples. Thereof, nine (45%) isolates were STEC O156:H25, three (15%) isolates typed STEC O43:H2. The remaining isolates occurred as single serotypes or were O non-typeable. Eleven (55%) of the isolates harboured *stx2a*, nine (45%) *eae*, and 14 (70%) *ehx*, respectively. WGS revealed the presence of *iss* in 16 (80%), *subAB* in four (20%) and *astA* in two (10%) of the isolates. Furthermore, *espA*, *tccP*, *nleA*, *nleB*, *tccP*, and *tir* were found exclusively among STEC O156:H25.

Eleven different sequence types (ST) were detected. The most prominent was ST300/ST5343, which comprised STEC O156:H25. All STEC isolates were pan susceptible. The results indicate that dromedary camels in Kenya may be reservoirs of STEC, including serotypes associated to disease in humans. STEC in camels may represent a health hazard for humans with close contact or to consumers of camel derived foodstuffs.

Keywords: camels, STEC, serotypes, virulence genes, whole genome sequencing

Zusammenfassung

Shiga toxin-produzierende *Escherichia coli* (STEC) verursachen gastrointestinale Krankheiten. Um das Auftreten von STEC in kenianischen Kamelen zu untersuchen, wurden *E. coli* von 163 Kamelen auf *stx1* und *stx2* getestet. STEC Stämme wurden isoliert und serotypisiert. Mittels PCR wurden die *stx*-Gene subtypisiert und *eae* und *ehx* nachgewiesen. Whole genome sequencing (WGS) wurde durchgeführt, um weitere Virulenzgene zu detektieren und die Multilocus-Sequenztypen zu definieren. Antimikrobielle Resistenzprofile wurden mittels Disk Diffusion ermittelt.

STEC wurden aus 20 (12.3%) der Proben isoliert. Davon waren neun (45%) STEC O156:H25, drei (15%) STEC O43:H2. Die übrigen Isolate waren Einzelfälle oder O-untypisierbar. Elf (55%) Isolate borgen *stx2a*, neun (45%) *eae*, und 14 (70%) *ehx*. WGS ermittelte die Präsenz von *iss* in 16 (80%), *subAB* in vier (20%) und *astA* in zwei (10%) Isolaten. Weiter wurden *espA*, *tccP*, *nleA*, *nleB*, *tccP*, und *tir* nur in den zu STEC O156:H25 gehörenden Isolaten detektiert.

Elf verschiedene Sequenztypen wurden detektiert. Der häufigste war ST300/ST5343, welcher STEC O156:H25 enthält. Alle STEC Isolate waren pansensibel. Diese Resultate lassen darauf schliessen, dass die kenianischen Kamele potentielle STEC-Reservoirs sind. Einschliesslich potentiell humanpathogener Serotypen. STEC in Kamelen könnten ein Gesundheitsrisiko für Menschen mit engem Kontakt oder Konsumenten von kamelstämmigen Nahrungsmitteln sein.

Schlüsselwörter: Kamele, STEC, Serotypen, Virulenzgene, whole genome sequencing

Shiga toxin-producing *Escherichia coli* (STEC) isolated from fecal samples of African dromedary camels

Melinda Baschera¹, Nicole Cernela¹, Marc J.A. Stevens¹, Anne Liljander², Jörg Jores³, Victor Max Corman^{4,5}, Magdalena Nüesch-Inderbinen^{1*}, Roger Stephan¹

¹Institute for Food Safety and Hygiene, Vetsuisse Faculty, University of Zurich, Switzerland

²International Livestock Research Institute, PO Box 30709, 00100 Nairobi, Kenya

³Institute for Veterinary Bacteriology, Vetsuisse Faculty, University of Bern, Switzerland

⁴Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, and Berlin Institute of Health, Institute of Virology, Berlin

⁵German Centre for Infection Research, associated partner Charité, Berlin, Germany.

Keywords: camels, STEC, serotypes, virulence genes, whole genome sequencing

*Corresponding author: Magdalena Nüesch-Inderbinen, Institute for Food Safety and Hygiene, Vetsuisse Faculty, University of Zurich, Zurich, Switzerland.

Phone +41 44 635 86 51

E-mail: magdalena.nueesch-inderbinen@uzh.ch

Abstract

Shiga toxin producing *Escherichia coli* (STEC) cause gastrointestinal illnesses including non-bloody or bloody diarrhoea, haemorrhagic colitis (HC), and the haemolytic uremic syndrome (HUS). To investigate the occurrence of STEC among grazing dromedaries from Kenya, *E. coli* isolated from fecal matter collected from 163 dromedaries on a large ranch were screened for the presence of *stx1* and *stx2*. STEC strains were isolated and serotyped. Isolates were subjected to PCR for the subtyping of *stx* genes and for the detection of *eae* and *ehx*. In addition, whole genome sequencing (WGS) was carried out to detect further virulence genes and to determine the multilocus sequence types (MLST). Antimicrobial resistance profiles were determined by disk diffusion.

STEC was isolated from 20 (12.3%) of the fecal samples. Thereof, nine (45%) isolates were STEC O156:H25, three (15%) isolates typed STEC O43:H2. The remaining isolates occurred as single serotypes or were O non-typeable. Eleven (55%) of the isolates harboured *stx2a*, nine (45%) *eae*, and 14 (70%) *ehx*, respectively. WGS revealed the presence of *iss* in 16 (80%), *subAB* in four (20%) and *astA* in two (10%) of the isolates. Furthermore, *espA*, *tccP*, *nleA*, *nleB*, *tccP*, and *tir* were found exclusively among STEC O156:H25.

Eleven different sequence types (ST) were detected. The most prominent was ST300/ST5343, which comprised STEC O156:H25. All STEC isolates were pan susceptible to a panel of 16 antimicrobial agents. Overall, the results indicate that dromedary camels in Kenya may be reservoirs of STEC, including serotypes possessing virulence markers associated to disease in humans, such as STEC O156:H25. STEC in camels may represent a health hazard for humans with close contact to camels or to consumers of camel derived foodstuffs, such as unpasteurised camel milk.

Introduction

In East Africa, the one-humped Arabian camel (*Camelus dromedarius*) is an important livestock species particularly well adapted to semi-arid and arid environments [1]. In Kenya, the husbandry of semi-domesticated dromedaries is a traditional livelihood of the nomadic pastoralists, with milk constituting an important nutrient source and income asset [2].

Kenya as a country relies heavily on wildlife tourism [3]. In the Laikipia county in central Kenya which is known for its remarkable abundance of wildlife, livestock husbandry and tourism are alternatives to land cultivation [4, 5]. Dromedary camel herds have become established both in pastoral communities and on large ranches where they are used for wildlife safaris and to produce milk for consumption and trade [1].

Camels in sub-saharan Africa reportedly represent a reservoir for a variety of potentially zoonotic diseases and pathogens, including *Coxiella burnetii* [6], *Brucella* spp. [7], Hepatitis E Virus [8], and Middle East respiratory syndrome coronavirus [9]. By contrast, Shiga toxin producing *Escherichia coli* (STEC), which may cause gastrointestinal illnesses including non-bloody or bloody diarrhoea, haemorrhagic colitis (HC), and the haemolytic uremic syndrome (HUS) [10], has rarely been described in camels. STEC causes an estimated 2,801,000 acute illnesses annually and poses a significant health burden worldwide [11, 12].

The distinguishing feature of STEC is the presence of one or more Shiga toxin (*stx*) genes which include two major forms, *stx1* and *stx2* and subtypes thereof [13]. Other major virulence genes that influence the pathogenic potential of STEC include the intimin gene *eae*, and the enterohemolysin gene *ehx* [14, 15]. Additional virulence

factors such as *espA*, *nleA*, *iss* or *subAB* are also frequently associated with disease in humans [16, 17].

STEC O157:H7 is the serotype most frequently associated with severe clinical outcomes, nonetheless, non-O157 STEC serogroups, including O26, O103, O111, and O145, are also recognized for their pathogenic potential [18].

Wild and domestic animals, including cattle, sheep, goats and deer represent the most important reservoirs of STEC [12]. Examples of healthy domestic animals that are known to be hosts of human pathogenic STEC serotypes are given in Table 1.

The consumption of contaminated water, meat, raw milk, raw milk products and produce, and the direct contact with farm animals have been identified as important routes of transmission [12, 19]. A better understanding of the occurrence of STEC among animal reservoirs is essential to provide a framework for developing approaches to reduce the burden of STEC related disease worldwide.

So far, *E. coli* O157:H7 and STEC O157:H7 was isolated from meat and feces of camels in Iran, respectively [20, 21]. By contrast, several studies have reported the absence of STEC in camels in the United Arab Emirates and Eastern Africa [22, 23, 24].

This study aimed to update the current understanding on the occurrence of STEC in camels in Eastern Africa by screening fecal samples of a herd of healthy animals kept under extensive husbandry systems on a ranch in Laikipia, Kenya, and to characterize all STEC strains with regard to their serotypes and their virulence potential.

Material and Methods

Sample collection. Fecal samples of 163 healthy dromedary camels (117 females, 45 males and one of unknown gender) located in a ranch in the northern Laikipia county in Kenya were collected during January and February 2017. The camels were extensively kept, they shared their pasture with game and cattle. Fecal matter was collected by rectal palpation using one glove per animal. Samples were placed in sterile containers, cooled for transport and frozen at -80°C.

The sample collection was performed in strict accordance with the Kenyan legislation for animal experimentation and were approved by the institutional animal care and use committee (IACUC reference number 2014.08). Since 1993, the International Livestock Research Institute (ILRI) in Kenya has complied voluntarily with the United Kingdom's Animals (Scientific Procedures) Act 1986 that contains guidelines and codes of practice for the housing and care of animals used in scientific procedures.

Screening for *stx* by PCR. Each fecal sample was enriched at a 1:10 ratio in Enterobacteriaceae enrichment (EE) broth (Becton, Dickinson, Heidelberg, Germany) for 24 h at 37°C. One loopful of each of the enrichment cultures was cultured on sheep blood agar (Difco™ Columbia Blood Agar Base EH; Becton Dickinson AG, Allschwil, Switzerland) using the streak plate technique. The resulting colonies were washed off with 2 ml 0.85% NaCl and DNA was extracted by a standard lysis protocol. Screening for *stx1* and *stx2* genes was performed by real-time PCR (LightCycler R 2.0 Instrument, Roche Diagnostics Corporation, Indianapolis, IN, USA) using the QuantiFast Multiplex PCR Kit (Qiagen, Hombrechtikon, Switzerland) according to the guidelines of the European Union Reference Laboratory (EURL) [25].

Recovery of STEC. In the event of a *stx* positive PCR result, one loopful each of the enrichment broth was streaked onto on sheep blood agar (Difco™ Columbia Blood Agar Base EH; Becton Dickinson AG, Allschwil, Switzerland) and onto RAPID' *E. coli* Agar (BioRad, Basel, Switzerland) to obtain single colonies of *E. coli*.

From each plate, individual colonies were picked and suspended in 0.5 ml NaCl 0.85%. Isolates that were confirmed to possess *stx* (*stx1* and/or *stx2*) by real-time PCR were subcultured on sheep blood agar and the presence of *stx* was confirmed using the Assurance GDS® for Shiga Toxin Genes (Bio Control Systems, Bellevue, WA, USA). From plates yielding more than one *stx* positive colony, one isolate was randomly chosen for subsequent characterization.

Serotyping and molecular typing of *stx*, *eae* and *ehx* genes. Isolates were serotyped using the *E. coli* SeroGenoTyping AS-1 Kit (Alere Technologies, Jena, Germany). The determination of *stx1* subtypes (*stx1a*, *stx1c*, *stx1d*) and *stx2* subtypes (*stx2a*, *stx2b*, *stx2c*, *stx2d*, *stx2e*, *stx2g*) was performed by conventional PCR amplification [13]. Screening for *eae*, and *ehx* was performed by real-time PCR according to the guidelines of the EURL [26].

Whole genome sequencing. Genomic DNA libraries were prepared using the Nextera™ DNA Flex Library Preparation Kit following the manufacturer's protocol (Illumina, San Diego, CA, USA). Libraries were loaded on a 300-cycle, 2 × 150 bp paired-end read cartridge (Illumina MiniSeq High Output Reagent Kit) and run on a MiniSeq Illumina sequencer (Illumina). The Spades assembler v3.12 [27] was used to assemble the reads with using the option "--careful". The spades output was filtered for size > 1000 bp and coverage > 25-fold resulting in the final assemblies that were submitted to NCBI GenBank.

The sequence reads were used to run analysis in the Advanced Research Infrastructure for Experimentation in genomics (ARIES) public Galaxy server (<https://w3.iss.it/site/aries/>), i.e. quality control, assembly, virulence gene analysis, Shiga-toxin analysis, and serotyping.

To assess the genetic relatedness of the isolates, core genome multilocus sequence typing (cgMLST) was carried out using Ridom Seq-Sphere Software (version 4.1.9, available at <http://www.ridom.de/seqsphere/cgmlst/>). The MLST schemes “Warwick” and “Pasteur” were used for MLST-typing, and the Enterobase *Escherichia/Shigella* v1 scheme (<http://enterobase.warwick.ac.uk>) for cgMLST typing. Assembled genomes were blasted against the reference database using standard settings and the option “ignore missing values pairwise” and “Discard genomes with > 3% missing genes” were used for typing. Trees were visualised using the minimal spanning tree option in seqsphere.

Antibiotic resistance profiles. Antimicrobial susceptibility testing was performed using the disk-diffusion method and the antibiotics ampicillin (AM), amoxicillin-clavulanic acid (AMC), cefazolin (CZ), cefotaxime (CTX), cefepime (FEP), nalidixic acid (NA), ciprofloxacin (CIP), gentamicin (GM), kanamycin (K), streptomycin (S), sulfamethoxazole/trimethoprim (SXT), fosfomycin (FOS), azithromycin (AZM), nitrofurantoin (F/M), chloramphenicol (C) and tetracycline (T) (Becton Dickinson, Heidelberg, Germany). Results were interpreted according to Clinical and Laboratory Standards Institute (CLSI) performance standards [28].

Results and discussion

In sub-Saharan Africa, little information is available on the epidemiology of STEC in humans, food and animals, and current knowledge of STEC sources needs to be improved [12]. Data on the occurrence and the characteristics of STEC in African camels is limited. This study provides evidence for the fecal carriage of STEC among dromedary camels located on a ranch in Laikipia Kenya. The restriction to one farm in Kenya is a limitation of this study and may have led to bias associated with camels from one herd only. In order to obtain a wider picture, analyses of feces from camels on other farms and other regions in Kenya would be required. For this investigation, a total of 163 fecal swabs were obtained. Thereof, 53 (32.5%) tested positive for *stx* by real-time PCR. STEC was recovered from 20 of the *stx* positive samples, amounting to 12.3% of all fecal samples. Overall, fecal carriage of STEC was confirmed for 14.5% (17 of a total of 117) of the female camels and 6.7% (3 of 45) of the male camels (Table 2).

The serotypes and virulence genotypes of the STEC isolates are summarized in Table 2. The most frequent serotypes included nine (45% of the isolates) STEC O156:H25 isolates and three (15%) O43:H2 isolates. The remaining strains occurred as single serotypes or were O non-typeable (Table 2). STEC O156:H25 has been described as a persistent colonizer of the bovine gut [29] and has also been linked to human infection in Germany and Switzerland [30, 31] (Table 1). By contrast, STEC O43:H2 is a rare serotype, but has been detected in beef in West Africa [32].

Further, various *stx* subtypes were detected among the isolates, including *stx2a* which is associated with severe disease in STEC infected humans [33]. The *stx2a* gene was found in a total of 11 (55%) of the isolates, including the nine STEC

O156:H25 strains, and the STEC O8:H49 and STEC Ont:H7 isolates, respectively (Table 2). The intimin gene *eae* was detected in nine (45%) isolates, all belonging to serotype O156:H25 (Table 2). While *eae* is one of the most prominent virulence factors contributing to pathogenesis, many STEC feature other virulence genes that may enhance their pathogenic potential, such as enterohaemorrhagic *E. coli* haemolysin *ehx* [15, 30]. In this study, *ehx* was detected by PCR in fourteen (70%) of the isolates (Table 2).

Furthermore, we used WGS to provide additional information on the virulence of the analysed STEC. Table 2 shows the presence or absence of a selection of virulence genes that predict severe disease in humans, including *espA*, *tccP*, *nleA*, *nleB* and *tir* (de Boer et al., 2015). Other virulence factors included in Table 2 are *iss*, which encodes an increased serum survival protein and is associated with non-bloody diarrhoea as well as extra-intestinal pathogenic infections [34], *subAB*, which is an emerging pathogenic factor among *eae* negative human pathogenic STEC isolates [17], and *astA* encoding a heat-stable enterotoxin EAST1 that is found mostly in enterohaemorrhagic and enteroaggregative *E. coli* [35]. The complete ARIES analysis reports on the virulotypes of the 20 sequenced strains are available upon request.

Taken together, *espA*, *tccP*, *nleA*, *nleB* and *tir* were detected exclusively among the nine STEC O156:H25 isolates, while *iss* was identified in 16 (80%), and *subAB* in four (20%) of the isolates, respectively (Table 2). A minority of two strains (10%) possessed *astA* (Table 2). Although all STEC can conceivably cause diarrhoea, the risk and severity of infection is linked to the presence of virulence factors, in particular *stx2a* and *eae* [12]. Our results show that although 45% of the STEC isolated from camel feces are not highly pathogenic according to their virulence

factor profiles, the majority (55%) is associated with *stx2a*, and STEC O156:H25 isolates in particular, exhibit a virulotype that suggests a potential health risk. Using the Pasteur scheme, four different sequence types (ST) were identified among 14 isolates (Table 2). Using the Warwick scheme, 11 different STs were identified (Table 2). Of the nine STEC O156:H25 isolates, eight grouped in the sequence type ST300 and ST5334, which differ by one nucleotide in the *gyrB* gene, indicating close phylogenetic relationship of the isolates. Similarly, the three STEC O43:H2 isolates were assigned to ST937 and ST6275 which differ by one nucleotide in *fumC*. Two STEC Ont:H20 possessed ST5759. All other isolates belonged to singly occurring STs (Table 2). Accordingly, cgMLST using Ridom SeqSphere+ software divided the strains into distinct clusters that comprised the STEC O156:H25 isolates, the STEC O43:H2 isolates, and the Ont:H20 isolates, respectively, while the remaining strains did not reveal close phylogenetic relationship (Figure 1). Finally, all STEC isolates remained fully susceptible to all antimicrobials tested (data not shown).

Conclusions

The data from our study suggest that dromedary camels are a reservoir of human pathogenic STEC. While none of the fecal samples contained STEC O157:H7 or other non-O157 STEC serogroups frequently associated with severe clinical outcomes such as bloody diarrhoea, HC, or HUS, the majority of the isolates harboured *stx2a*, and the STEC O156:H25 isolates harboured *eae*, *ehx*, and other virulence factors often associated with disease in humans. Their occurrence in fecal samples from camels may represent a threat to humans, especially to those who live in long-term close contact with camels and consume raw camel milk.

Nucleotide sequences

The nucleotide sequences have been deposited at GenBank under accession nos. **RJDB000000000-RJBU000000000.**

Disclosures

The authors report no conflict of interest. All authors have read and approved the final article.

Acknowledgements

This work was partly supported by the University of Zurich and the Swiss Federal Office of Public Health, Division communicable Disease. Additional funds were received from the German ministry of Economic Cooperation (Contract No.

81195004). Anne Liljander was supported by the Centre for International Migration and Development. We thank Katrin Zurfluh for technical assistance.

References

- [1] Schwartz, H.J., M. Dioli. The one-humped camel (*Camelus dromedarius*) in eastern Africa. A pictorial guide to diseases, health care, and management. 1st ed. Berlin, 1992.
- [2] Benka, V. Human-wildlife conflict, interspecies disease, and justice in a wild-life rich region in Kenya. University of Michigan. Master's thesis. Available at <https://deepblue.lib.umich.edu/bitstream/handle/2027.42/96190/benka%20thesis%20final.pdf?sequence=1>. Accessed 29 Nov 2018.
- [3] Valle, E., M.N. Yobesia. Economic contribution of tourism in Kenya. *Tourism Analysis*. 2009;14:401-4.
- [4] Kinnaird, M.F., T.G. O'brien. Effects of private-land use, livestock management, and human tolerance on diversity, distribution, and abundance of large African mammals. *Conserv Biol*. 2012;26:1026-39.
- [5] Sundaresan, S.R., C. Riginos. Lessons learned from biodiversity conservation in the private lands of Laikipia, Kenya. *Great Plains Res.* (2010). 17-27. Available at <https://www.jstor.org/stable/23782172>. Accessed 28 Nov 2018.
- [6] DePuy, W., V. Benka, A. Massey, S.L. Deem, M. Kinnaird, T. O'Brien *et al.* Q fever risk across a dynamic, heterogeneous landscape in Laikipia County, Kenya. *EcoHealth*.. 2014;11:429-33.
- [7] Ducrotoy, M., W.J. Bertu, G. Matope, S. Cadmus, R. Conde-Álvarez, A.M. Gusi *et al.* Brucellosis in sub-saharan Africa: current challenges for management, diagnosis and control. *Acta Trop*. 2017;165:179-93.
- [8] Rasche, A., M. Saqib, A.M. Liljander, S. Bornstein, A. Zohaib, S. Renneker *et al.* Hepatitis E virus infection in dromedaries, north and east Africa, United Arab Emirates, and Pakistan, 1983-2015. *Emerg Infect Dis*. 2016;22:1249-52.

- [9] Jores, J. Middle East respiratory syndrome-coronavirus in camels: an overview for Sub-Saharan and North Africa. Evidence on Demand, UK. 2015.
DOI=http://dx.doi.org/10.12774/eod_cr.july2015.joresj.
- [10] Karch, H., P.I. Tarr, M. Bielaszewska. Enterohaemorrhagic *Escherichia coli* in human medicine. Int J Med Microbiol. 2005;295:405-418.
DOI=10.1016/j.ijmm.2005.06.009.
- [11] Majowicz, S. E., Scallan, E., Jones-Bitton, A., Sargeant, J. M., Stapleton, J., Angulo, F. J. *et al.* Global incidence of human Shiga toxin–producing *Escherichia coli* infections and deaths: A systematic review and knowledge synthesis. Foodborne Pathog. Dis. 2014;11:447-55.
DOI=10.1089/fpd.2013.1704
- [12] FAO (Food and Agriculture Organization of the United Nations), WHO (World Health Organization). Shiga Toxin-Producing *Escherichia coli* (STEC) and food: Attribution, characterization, and monitoring: Report. FAO; Rome, WHO; Geneva, 2018.
- [13] Scheutz, F., L.D. Teel, L. Beutin, D. Piérard, G. Buvens, H. Karch *et al.* Multicenter evaluation of a sequence-based protocol for subtyping Shiga toxins and standardizing Stx nomenclature. J Clin Microbiol. 2012;50:2951-63.
DOI=10.1128/JCM.00860-12.
- [14] Kaper, J.B., J.P. Nataro, H.L. Mobley. Pathogenic *Escherichia coli*. Nat Rev Microbiol. 2004;2:123-140. DOI=10.1038/nrmicro818.
- [15] Schmidt, H., L. Beutin, H. Karch. Molecular analysis of the plasmid-encoded hemolysin of *Escherichia coli* O157:H7 strain EDL 933. Infect Immun. 1995;63:1055-61.
- [16] de Boer, R.F., M. Ferdous, A. Ott, H.R. Scheper, G.J. Wisselink, M.E. Heck *et*

- al.* Assessing the public health risk of Shiga-toxin producing *Escherichia coli* using a rapid diagnostic screening algorithm. J Clin Microbiol. 2015 53:1588 – 98. DOI=10.1128/JCM.03590-14.
- [17] Paton, A.W., P. Srimanote, U.M. Talbot, H. Wang, J.C. Paton. A new family of potent AB(5) cytotoxins produced by Shiga toxigenic *Escherichia coli*. J Exp Med. 2004;200:35-46.
- [18] EFSA (European Food Safety Authority). The European Union summary report on trends and sources of zoonoses, zoonotic agents and food-borne outbreaks in 2016. EFSA Journal (2017);15(12):5077, 228 pp.
- [19] Karmali, M. A. Factors in the emergence of serious human infections associated with highly pathogenic strains of Shiga toxin-producing *Escherichia coli*. Int J Med Microbiol. 2018; 308:1067-1072. DOI= 10.1016/j.ijmm.2018.08.005.
- [20] Hajian, S., E. Rahimi, H. Mommtaz. A 3-year study of *Escherichia coli* O157: H7 in cattle, camel, sheep, goat, chicken and beef minced meat. Proc Internat Conf Food Eng Biotechnol (IPCBE'11); 2011; Singapore. IACSIT Press.
- [21] Salehi, T.Z., A. Tonelli, A. Mazza, H. Staji, P. Badagliacca, I.A. Tamai *et al.* Genetic characterization of *Escherichia coli* O157: H7 strains isolated from the one-humped camel (*Camelus dromedarius*) by using microarray DNA technology. Mol Biotechnol. 2012;51:283-88.
- [22] El-Sayed, A., S. Ahmed, W. Awad. Do camels (*Camelus dromedarius*) play an epidemiological role in the spread of Shiga toxin producing *Escherichia coli* (STEC) infection? Trop Anim Health Prod. 2008;40:469-73.
- [23] Moore, J.E., M. McCalmont, J. Xu, G. Nation, A.H. Tinson, L. Crothers *et al.* P Prevalence of faecal pathogens in calves of racing camels (*Camelus*

- dromedarius*) in the United Arab Emirates. Trop Anim Health Prod. 2002;34:283-87.
- [24] Tabatabaei, S., T.Z. Salehi, M.A. Badouei, I.A. Tamai, V. Akbarinejad, R. Kazempoor *et al.* Prevalence of Shiga toxin-producing and enteropathogenic *Escherichia coli* in slaughtered camels in Iran. Small Rumin Res. 2013;113:297-300.
- [25] EURL (European Union Reference Laboratory). 2013a. Identification and characterization of Verocytotoxin-producing *Escherichia coli* (VTEC) by PCR amplification of the main virulence genes. EU-RL VTEC_Method_01_Rev_0. Available online at: <http://old.iss.it/vtec/index.php?lang=2&tipo=3>. Accessed 28 Nov 2018.
- [26] EURL (European Union Reference Laboratory). 2013b. Identification and characterization of Verocytotoxin-producing *Escherichia coli* (VTEC) by real time PCR amplification of the main virulence genes and the genes associated with the serogroups mainly associated with severe human infections. EU-RL VTEC_Method_02_Rev 0. Available online at: <http://old.iss.it/vtec/index.php?lang=2&tipo=3>. Accessed 28 Nov 2018.
- [27] Bankevich, A., S. Nurk, D. Antipov, A.A. Gurevich, M. Dvorkin, A.S. Kulikov *et al.* SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. J Comput Biol. 2012;19:455-77.
- [28] CLSI (Clinical and Laboratory Standards Institute). Performance standards for antimicrobial susceptibility testing. 27th ed. CLSI supplement M100S. Wayne, 2017.
- [29] Barth, S.A., C. Menge, I. Eichhorn, T. Semmler, L.H. Wieler, D. Pickard *et al.* The accessory genome of Shiga toxin-producing *Escherichia coli* defines a

- persistent colonization type in cattle. *Appl Environ Microbiol.* 2016;82:5455-64.
- [30] Beutin, L., G. Krause, S. Zimmermann, S. Kaulfuss, K. Gleier. Characterization of Shiga toxin-producing *Escherichia coli* strains isolated from human patients in Germany over a 3-year period. *J Clin Microbiol.* 2004;42:1099-1108.
- [31] Fierz, L., N. Cernela, E. Hauser, M. Nüesch-Inderbinen, R. Stephan. Characteristics of Shigatoxin-producing *Escherichia coli* strains isolated during 2010-2014 from human infections in Switzerland. *Front Microbiol.* 2017;8:1471. DOI=10.3389/fmicb.2017.01471.
- [32] Martikainen, O., A. Kagambega, I.J. Bonkougou, N. Barro, A. Siitonen, K. Haukka. Characterization of shigatoxigenic *Escherichia coli* strains from Burkina Faso. *Foodborne Path Dis.* 2012;9:1015-21.
- [33] Fuller, C.A., C.A. Pellino, M.J. Flagler, J.E. Strasser, A.A. Weiss. Shiga toxin subtypes display dramatic differences in potency. *Infect Immun.* 2011;79:1329-37.
- [34] Matussek, A., C. Jernberg, I.M. Einemo, S. Monecke, R. Ehricht, I. Engelmann *et al.* Genetic makeup of Shiga toxin-producing *Escherichia coli* in relation to clinical symptoms and duration of shedding: a microarray analysis of isolates from Swedish children. *Eur J Clin Microbiol Infect Dis.* 2017;36:1433-41.
- [35] Maluta, R.P., J.L. Leite, T.C.G. Rojas, I.C.A. Scaletsky, E.A.L. Guastalli, M.D.C. Ramos *et al.* Variants of *astA* gene among extra-intestinal *Escherichia coli* of human and avian origin. *FEMS Microbiol Lett.* 2017;3646, fnw285. DOI=10.1093/femsle/fnw285.
- [36] Doane, C. A., Pangloli, P., Richards, H. A., Mount, J. R., Golden, D. A., Draughon, F. A. Occurrence of *Escherichia coli* O157:H7 in diverse farm environments. *J Food Prot.* 2007: 70: 6-10.

- [37] Persad, A. K. LeJeune, J. T. Animal reservoirs of Shiga toxin-producing *Escherichia coli*. Microbiol Spectr. 2014;2:231-44.
DOI=10.1128/microbiolspec.EHEC-0027-2014.
- [38] Kang, E., Hwang, S. Y., Kwon, K. H., Kim, K. Y., Kim, J. H., Park, Y. H. Prevalence and characteristics of Shiga toxin-producing *Escherichia coli* (STEC) from cattle in Korea between 2010 and 2011. J Vet Sci. 2014;15: 369-79. DOI=10.4142/jvs.2014.15.3.369.
- [39] Mainga, A. O., Cenci-Goga, B. T., Malahlela, M. N., Tshuma, T., Kalake, A., Karama, M. Occurrence and characterization of seven major Shiga toxin-producing *Escherichia coli* serotypes from healthy cattle on cow–calf operations in South Africa. Zoonoses Public Health. 2018;65:777-89.
DOI=10.1111/zph.12491.
- [40] Beutin, L., Geier, D., Steinrück, H., Zimmermann, S., Scheutz, F. Prevalence and some properties of Verotoxin (Shiga-like toxin)-producing *Escherichia coli* in seven different species of healthy domestic animals. J Clin Microbiol. 1993; 31: 2483-8. DOI=0095-1137/93/092483-06\$02.00/0.
- [41] Orden, J. A., Domínguez-Bernal, G., de la Fuente, R., Carrión, J. Subtilase cytotoxin-encoding *subAB2* variants in Verotoxin-producing *Escherichia coli* strains isolated from goats and sheep. Res Vet Sci. 2016;105: 74-6. DOI=10.1016/j.rvsc.2016.01.017.
- [42] Cortés, C., De la Fuente, R., Blanco, J., Blanco, M., Blanco, J. E., Dhahi, G. *et al.* Serotypes, virulence genes and intimin types of Verotoxin-producing *Escherichia coli* and enteropathogenic *E. coli* isolated from healthy dairy goats in Spain. Vet Microbiol. 2005;110: 67-76. DOI=10.1016/j.vetmic.2005.06.009.

- [43] Brandal, L. T., Sekse, C., Lindstedt, B.-A., Sunde, M., Løbersli, I., Urdahl, A. M.
et al. Norwegian sheep are an important reservoir for human-pathogenic
Escherichia coli O26:H11. Appl Environ Microbiol. 2012;78:4083-91.
DOI=10.1128/AEM.00186-12

Table 1: Examples of healthy (semi)-domestic animals as hosts of Shiga toxin producing *Escherichia coli* (STEC) serotypes associated with disease in humans.

Animal	STEC serotype(s)	Reference(s)
Camels	O157:H7, O156:H25	[21], this study
Cattle	O157:H7, O156:H25, O8:H19, O26:H2, O26:H8, O26:H11, O26:H19, O26:H21, O45:H2, O103:H2, O103:H21, O121:H8, O145:H2, O145:H8, O145:H28, O146:H21	[29, 36, 37, 38, 39, 40]
Chicken	O157:H7	[36, 37]
Goats	O146:H21, O82:H8, O128:H2	[40, 41, 42]
Sheep	O157:H7, O26:H11, O146:H21, O128:H2	[37, 40, 41, 43]
Swine	O157:H7	[36, 37]
Turkeys	O157:H7	[36]

Table 2: Characteristics of Shiga toxin producing *Escherichia coli* (STEC) isolated from feces of camels from Kenya.

Strain ID	Host gender	Serotype	Virulence genes											MLST		
			<i>stx</i> subtype	<i>eae</i>	<i>ehx</i>	<i>astA</i>	<i>espA</i>	<i>iss</i>	<i>nleA</i>	<i>nleB</i>	<i>tccP</i>	<i>tir</i>	<i>subAB</i>	ST Warwick	ST Pasteur	GenBank accession no.
C59	F	O8:H19	<i>stx1a</i>	-	+	-	-	+	-	-	-	-	-	201	294	RJDD000000000
C102	F	O8:H49	<i>stx2a</i>	-	+	-	-	+	-	-	-	-	-	111	nd	RJDT000000000
C139	F	O43:H2	<i>stx1c</i>	-	-	-	-	+	-	-	-	-	-	937	363	RJDN000000000
C119	F	O43:H2	<i>stx1a</i>	-	-	-	-	+	-	-	-	-	+	6275	363	RJDS000000000
C160	F	O43:H2	<i>stx1c</i>	-	-	-	-	+	-	-	-	-	+	6275	363	RJDK000000000
C34	F	O44:H18	<i>stx1a</i> , <i>stx2c</i>	-	+	-	-	+	-	-	-	-	+	6507	nd	RJDG000000000
C21	F	O117:H12	<i>stx1a</i>	-	-	-	-	+	-	-	-	-	-	101	nd	RJDJ000000000
C22	F	O150:H8	<i>stx1c</i>	-	+	+	-	-	-	-	-	-	+	906	303	RJDI000000000
C12	F	O156:H25	<i>stx2a</i>	+	+	-	+	+	+	+	+	-	-	300	591	RJDR000000000
C15	F	O156:H25	<i>stx2a</i>	+	+	-	+	+	+	+	-	+	-	300	591	RJDM000000000
C16	F	O156:H25	<i>stx2a</i>	+	+	-	-	-	-	-	-	-	-	300	591	RJDL000000000
C25	F	O156:H25	<i>stx2a</i>	+	+	-	+	+	+	+	+	+	-	300	591	RJDH000000000
C43	F	O156:H25	<i>stx2a</i>	+	+	-	+	-	+	+	-	+	-	300	591	RJDF000000000
C62	F	O156:H25	<i>stx2a</i>	+	+	-	+	+	+	+	-	+	-	300	591	RJDC000000000
C85	F	O156:H25	<i>stx2a</i>	+	+	-	+	+	+	+	-	+	-	300	591	RJDB000000000
C129	M	O156:H25	<i>stx2a</i>	+	+	-	+	+	+	+	-	+	-	300	591	RJDP000000000
C101	M	O156:H25	<i>stx2a</i>	+	+	+	+	+	+	+	+	+	-	5343	591	RJDU000000000
C127	F	Ont:H7	<i>stx1a</i> , <i>stx2a</i>	-	+	-	-	-	-	-	-	-	-	1308	nd	RJDQ000000000
C56	M	Ont:H20	<i>stx1c</i>	-	-	-	-	+	-	-	-	-	-	5759	nd	RJDE000000000
C131	F	Ont:H20	<i>stx1c</i>	-	-	-	-	+	-	-	-	-	-	5759	nd	RJDO000000000

astA, gene encoding enteroaggregative heat-stable toxin; *eae*, intimin gene; *ehx*, enterohemolysin gene; *espA*, gene encoding LEE-associated secreted translocator protein; *iss*, gene encoding increased serum survival protein; MLST, multilocus sequence type; nd, not defined; *nle*, genes encoding non-LEE-encoded factors; ST, sequence type; *subAB*, subtilase cytotoxin gene; *tccP*, gene encoding Tir cytoskeleton coupling protein; *tir*, gene encoding translocated intimin receptor; +, the gene is present; –, the gene is absent

Figure 1

Phylogenetic relationship of 20 Shiga toxin producing *Escherichia coli* (STEC) isolated from camel feces based on their multilocus sequence type (MLST) allelic profiles. The minimum spanning tree was generated using SeqSphere (Ridom GmbH). Numbers in the lines indicate the number of allele differences between isolates. The size of each circle reflects the number of isolates allocated to an individual sequence type (ST). The colours of the circles represent STs according to the Warwick scheme (<http://enterobase.warwick.ac.uk>). Strain IDs are indicated in the circles.

Danksagung

Für die Unterstützung während der Forschung, des Schreibens und der Fertigstellung dieser Dissertation danke ich meiner Familie, meinen Freunden und dem gesamten Team am Institut für Lebensmittelsicherheit und –hygiene der Vetsuisse Fakultät Zürich.

Curriculum Vitae

Vorname Name	Melinda Baschera
Geburtsdatum	27.04.1994
Geburtsort	Bülach, Zürich
Nationalität	Schweizerin
Heimatort	Zürich
08/2000 – 06/2006	Primarschule , Volketswil ZH, Schweiz
08/2006 – 06/2012	Kantonale Mittelschule , Zürich ZH, Schweiz
02.07.2012	Matura , Kantonale Mittelschule, Zürich ZH, Schweiz
09/2012 – 01/2017	Studium der Veterinärmedizin , Universität Zürich, Zürich, Schweiz
29. 12. 2017	Staatsexamen vet. med. , Universität Zürich, Zürich, Schweiz
02/2018 – 10/2020	Anfertigung der Dissertation unter Leitung von Prof. Dr. med. vet. Dr. h.c Roger Stephan am Institut für Lebensmittelsicherheit und –hygiene der Vetsuisse-Fakultät Universität Zürich Direktor: Prof. Dr. med. vet. Dr. h.c Roger Stephan